

TESTOSTERONE

MEDICATION(S) SUBJECT TO STEP THERAPY

ANDRODERM, DEPO-TESTOSTERONE, FORTESTA, METHYLTESTOSTERONE 10 MG CAP, NATESTO, STRIANT, TESTONE CIK, TESTOPEL, TESTOSTERONE 100 MG PELLET, TESTOSTERONE 12.5 MG/ACT (1%) GEL, TESTOSTERONE 200 MG PELLET, TESTOSTERONE 50 MG PELLET, TESTOSTERONE ENANTHATE 200 MG/ML SOLUTION, VOGELXO, VOGELXO PUMP

CRITERIA

Patient must have previously tried generic testosterone cypionate and one other generic testosterone product

2717: DMT2 GLP1

MEDICATION(S) SUBJECT TO STEP THERAPY

BYDUREON, BYDUREON BCISE, BYETTA 5 MCG PEN, OZEMPIC (0.25 OR 0.5 MG/DOSE), OZEMPIC (1 MG/DOSE), OZEMPIC (2 MG/DOSE), RYBELSUS, TRULICITY

CRITERIA

Patient must have previous trial of Metformin

2717: MOUNJARO

MEDICATION(S) SUBJECT TO STEP THERAPY

MOUNJARO

CRITERIA

Patient must have previously tried Metformin

HYPNOTICS - 1

MEDICATION(S) SUBJECT TO STEP THERAPY

EDLUAR, INTERMEZZO, ZOLPIMIST

CRITERIA

Patient must have previously tried zolpidem, zaleplon, or eszopiclone

ATTENTION-DEFICIT/HYPERACTIVITY DISORDER (ADHD)

MEDICATION(S) SUBJECT TO STEP THERAPY

DESOXYN, EVEKEO ODT, ZENZEDI, ADHANSIA XR, DAYTRANA, JORNAY PM, METHYLPHENIDATE HCL ER 18 MG TAB ER 24H, RITALIN, RITALIN LA

CRITERIA

Patient must have previously tried any 2 formulary generic CNS stimulant options

614: QUILLI

MEDICATION(S) SUBJECT TO STEP THERAPY

QUILLICHEW ER, QUILLIVANT XR

CRITERIA

Patient must have previously tried an extended release generic stimulant.

ANESTHETICS

MEDICATION(S) SUBJECT TO STEP THERAPY

NUVAKAAN-II

CRITERIA

Patient must have tried generic or individual components (lidocaine and prilocaine)

PSORIASIS

MEDICATION(S) SUBJECT TO STEP THERAPY

CLODERM, EUCRISA, PIMECROLIMUS, PROTOPIC 0.03 % OINTMENT, TACROLIMUS 0.03 % OINTMENT

CRITERIA

Patient must have tried a topical corticosteroid

IMMUNOSUPPRESSIVES

MEDICATION(S) SUBJECT TO STEP THERAPY

ASTAGRAF XL

CRITERIA

Patient must have previously tried generic tacrolimus.

CARDIOVASCULAR - 4

MEDICATION(S) SUBJECT TO STEP THERAPY

TEKTURNA HCT

CRITERIA

Patient must have tried generic ACE, ARB, ACE combination, or ARB combination

ANTIDEPRESSANTS

MEDICATION(S) SUBJECT TO STEP THERAPY

SAVELLA, SAVELLA TITRATION PACK

CRITERIA

Patient must have previous trial of a TCA (e.g. amitriptyline) or SNRI (e.g. duloxetine)

TOPICAL ANTIPARASITICS

MEDICATION(S) SUBJECT TO STEP THERAPY

LINDANE, SPINOSAD

CRITERIA

Patient must have previously tried permethrin

DERMATOLOGICALS

MEDICATION(S) SUBJECT TO STEP THERAPY

PICATO

CRITERIA

Patient must have previously tried topical fluorouracil or imiquimod

EPILEPSY - 2

MEDICATION(S) SUBJECT TO STEP THERAPY

QUDEXY XR, TROKENDI XR

CRITERIA

Patient must have previously tried generic topiramate IR

ANTIINCONTINENCE AGENT

MEDICATION(S) SUBJECT TO STEP THERAPY

GELNIQUE, OXYTROL

CRITERIA

Patient must have previously tried oxybutynin and tolterodine

GOUT: MITIGARE

MEDICATION(S) SUBJECT TO STEP THERAPY

MITIGARE

CRITERIA

Patient must have previously tried generic colchicine tablets

GOUT - 2

MEDICATION(S) SUBJECT TO STEP THERAPY

FEBUXOSTAT

CRITERIA

Patient must have previously tried allopurinol

ANTI-HISTAMINES - 2

MEDICATION(S) SUBJECT TO STEP THERAPY

DEXCHLORPHENIRAMINE MALEATE 2 MG/5ML SOLUTION, RYCLORA

CRITERIA

Patient must have tried diphenhydramine

ANTIMIGRAINE PREPARATIONS - 4

MEDICATION(S) SUBJECT TO STEP THERAPY

AMERGE, FROVA, SUMATRIPTAN SUCCINATE REFILL 6 MG/0.5ML SOLN CART, TOSYMRA, ZOLMITRIPTAN 2.5 MG SOLUTION, ZOLMITRIPTAN 5 MG SOLUTION, ZOMIG 2.5 MG SOLUTION, ZOMIG 5 MG SOLUTION

CRITERIA

Patient must have previously tried two preferred generic oral triptans

ANTIMIGRAINE PREPARATIONS - 2

MEDICATION(S) SUBJECT TO STEP THERAPY

TOSYMRA

CRITERIA

Patient must have previously tried two preferred generic oral triptans

ANTIPSYCHOTICS - 2

MEDICATION(S) SUBJECT TO STEP THERAPY

FANAPT, FANAPT TITRATION PACK

CRITERIA

Patient must have previously tried 2 of the following: olanzapine, quetiapine (IR/ER), risperidone/-ODT, aripiprazole

ANTIPSYCHOTICS - 4

MEDICATION(S) SUBJECT TO STEP THERAPY

PALIPERIDONE ER

CRITERIA

Patient must have previously tried 2 of the following: olanzapine, quetiapine (IR/ER), risperidone, aripiprazole.

HYPNOTICS - 3

MEDICATION(S) SUBJECT TO STEP THERAPY

RAMELTEON

CRITERIA

Patient must have previously tried eszopiclone, zaleplon, or zolpidem